

## NCTC REFERENCE CULTURES RECOMMENDED IN NATIONAL STANDARD OPERATING PROCEDURES (SOP's)

National Standard Methods have been developed since 1996 by working groups of experienced laboratory-based medical microbiologists and scientists. Each document has undergone a comprehensive consultation process to ensure that best possible practice is reflected in the methods. SOPs and guidelines that reflect such broad consensus promote practices of high quality, helping to assure the comparability of diagnostic information obtained in different laboratories and provide a reference point for method development. These procedures are intended solely as a general resource for practising professionals in the field operating in the UK and specialist advice should be obtained where necessary. For more information on the SOP's themselves, visit [www.evaluations-standards.org.uk](http://www.evaluations-standards.org.uk) or e-mail Standards@HPA.org.uk

### ORDER FORM

Customer details: Customer's Order Number: \_\_\_\_\_  
 Name \_\_\_\_\_ Title \_\_\_\_\_  
 Address \_\_\_\_\_  
 \_\_\_\_\_  
 Postcode \_\_\_\_\_ Telephone No. \_\_\_\_\_  
 Signature \_\_\_\_\_ Date \_\_\_\_\_

RETURN TO: NATIONAL COLLECTION OF TYPE CULTURES Certified to BS EN ISO  
 9001:2000 HPA Centre for Infections  
 61 Colindale Avenue  
 LONDON, NW9 5EQ  
 United Kingdom Tel: +44 (0)20 8870 2000 Fax: +44 (0)20 8205 7483

	<b>NCTC No.</b>	<b>Organisms for Media QC</b>	<b>MSOP number</b>	<b>Qty Req'd</b>
1	10215	<i>Actinomyces israelii</i>	25	
2	7464	<i>Bacillus cereus</i>	18	
3	10400	<i>Bacillus subtilis</i>	18	
4	9343	<i>Bacteroides fragilis</i>	3, 4, 25, 26	
5	10739	<i>Bordetella pertussis</i>	11	
6	10661	<i>Burkholderia cepacia</i>	9	
7	11322	<i>Campylobacter jejuni</i>	20	
8	NCPF 3089	<i>Candida albicans</i>	5, 6	
9	NCPF 3153	<i>Candida albicans</i>	6	
10	NCPF 3179	<i>Candida albicans</i>	5, 6, 20	
11	11204	<i>Clostridium difficile</i>	17	
12	8237	<i>Clostridium perfringens</i>	33, 36, 42	
13	532	<i>Clostridium sporogenes</i>	36	
14	10356	<i>Corynebacterium diphtheriae</i> var. <i>mitis</i> (non-toxigenic) *	7	
15	775	<i>Enterococcus faecalis</i>	2, 22, 39, 41, 44, 48, 50	
16	12697	<i>Enterococcus faecalis</i>	49	
17	9001	<i>Escherichia coli</i>	2, 6, 8, 10, 13, 14, 15, 16, 17, 18, 19, 20, 23, 24, 29, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 43, 44, 45, 46	
18	10418	<i>Escherichia coli</i>	49	
19	12900	<i>Escherichia coli</i> O157 VT-	23	
20	12241	<i>Escherichia coli</i>	49	
21	10945	<i>Haemophilus ducreyi</i>	28	

	<b>NCTC No.</b>	<b>Organisms for Media QC (Cont'd)</b>	<b>MSOP number</b>	<b>Qty Req'd</b>
22	8143	<i>Haemophilus influenzae</i>	1, 27	
23	11931	<i>Haemophilus influenzae</i>	1, 27	
24	12699	<i>Haemophilus influenzae</i>	49	
25	11637	<i>Helicobacter pylori</i>	24	
26	9528	<i>Klebsiella aerogenes</i>	31, 47	
27	662	<i>Lactococcus lactis</i>	41, 44	
28	12821	<i>Legionella pneumophila</i>	29	
29	11994	<i>Listeria monocytogenes</i>	22, 50	
30	8375	<i>Neisseria gonorrhoeae</i>	5	
31	12700	<i>Neisseria gonorrhoeae</i>	49	
32	10975	<i>Proteus mirabilis</i>	2, 5, 12, 37	
33	10662	<i>Pseudomonas aeruginosa</i>	8, 9, 35, 40, 49	
34	12934	<i>Pseudomonas aeruginosa</i>	49	
35	4840	<i>Salmonella poona</i>	13, 14, 19, 32, 38	
36	8574	<i>Shigella sonnei</i> *	13, 14	
37	6571	<i>Staphylococcus aureus</i>	1, 2, 5, 10, 11, 12, 21, 30, 34, 43, 45, 46, 48, 49	
38	8331	<i>Staphylococcus aureus</i>	30	
39	12493	<i>Staphylococcus aureus</i>	49	
40	12981	<i>Staphylococcus aureus</i>	49	
41	11047	<i>Staphylococcus epidermidis</i>	10, 21	
42	12977	<i>Streptococcus pneumoniae</i>	49	
43	8198	<i>Streptococcus pyogenes</i>	1, 12	
44	11218	<i>Vibrio furnissii</i>	16	
45	10885	<i>Vibrio parahaemolyticus</i>	16	
46	10460	<i>Yersinia enterocolitica</i>	15	
		<b>Strains for Food, Dairy &amp; Water Products</b>	<b>MSOP Number</b>	
47	5866	<i>Acinetobacter lwoffii</i>	W2	
48	8049	<i>Aeromonas hydrophila</i>	W9	
49	13216	<i>Escherichia coli</i>	F16	
2	7464	<i>Bacillus cereus</i>	F15	
3	10400	<i>Bacillus subtilis</i>	F15	
7	11322	<i>Campylobacter jejuni</i>	F21, W8	
12	8237	<i>Clostridium perfringens</i>	F14, W5	
15	775	<i>Enterococcus faecalis</i>	D2, D3, F19, W3	
17	9001	<i>Escherichia coli</i>	D2, D3, D4, D5, F13, F15, F16, F17, F20, F21, F23, W2, W3, W5, W6, W7, W8, W9, W10, W16, W18	
19	12900	<i>Escherichia coli</i> O157 VT-	F17, W16	
26	9528	<i>Klebsiella aerogenes</i>	D5, F16, F20, W2, W4, W18	
27	662	<i>Lactococcus lactis</i>	D2, D3	
29	11994	<i>Listeria monocytogenes</i>	F19	
33	10662	<i>Pseudomonas aeruginosa</i>	F23, W2, W6, W8, W18	
35	4840	<i>Salmonella poona</i>	F13, W7	
37	6571	<i>Staphylococcus aureus</i>	D4, F12, W10	
41	11047	<i>Staphylococcus epidermidis</i>	F12, W10	
44	11218	<i>Vibrio furnissii</i>	W9	

\* Hazardous pathogen, will only be supplied on receipt of an order signed by a designated member of staff for whom we have a specimen signature (see NCTC Conditions of Supply). All cultures are priced at £50.00 each, with a 10% discount for orders ≥ 10 and a 20% additional discount to UK Health and Educational Customers.

Note: VAT, where applicable, is additional (EU customers please state VAT No. \_\_\_\_\_).

Postage and packaging: within UK £10 per order; outside UK £90 - £160 per order depending on country.

May 2005

