

NATIONAL STANDARD METHOD

UNDER REVIEW

INVESTIGATION OF GASTRIC ASPIRATES AND INFECTION SCREEN SWABS FROM NEONATES

BSOP 23

Issued by Standards Unit, Evaluations and Standards Laboratory
Specialist and Reference Microbiology Division

Association of Medical Microbiologists
Association of Medical Microbiologists
Association of Medical Microbiologists



INVESTIGATION OF GASTRIC ASPIRATES AND INFECTION SCREEN SWABS FROM NEONATES

Issue no: 4.1 Issue date: 03.05.05 Issued by: Standards Unit, Evaluations and Standards Laboratory Page 1 of 19
Reference no: BSOP 23i4.1

This SOP should be used in conjunction with the series of other SOPs from the Health Protection Agency

www.evaluations-standards.org.uk

Email: standards@hpa.org.uk

STATUS OF NATIONAL STANDARD METHODS

National Standard Methods, which include standard operating procedures (SOPs), algorithms and guidance notes, promote high quality practices and help to assure the comparability of diagnostic information obtained in different laboratories. This in turn facilitates standardisation of surveillance underpinned by research, development and audit and promotes public health and patient confidence in their healthcare services. The methods are well referenced and represent a good minimum standard for clinical and public health microbiology. However, in using National Standard Methods, laboratories should take account of local requirements and may need to undertake additional investigations. The methods also provide a reference point for method development.

National Standard Methods are developed, reviewed and updated through an open and wide consultation process where the views of all participants are considered and the resulting documents reflect the majority agreement of contributors.

Representatives of several professional organisations, including those whose logos appear on the front cover, are members of the working groups which develop National Standard Methods. Inclusion of an organisation's logo on the front cover implies support for the objectives and process of preparing standard methods. The representatives participate in the development of the National Standard Methods but their views are not necessarily those of the entire organisation of which they are a member. The current list of participating organisations can be obtained by emailing standards@hpa.org.uk.

The performance of standard methods depends on the quality of reagents, equipment, commercial and in-house test procedures. Laboratories should ensure that these have been validated and shown to be fit for purpose. Internal and external quality assurance procedures should also be in place.

Whereas every care has been taken in the preparation of this publication, the Health Protection Agency or any supporting organisation cannot be responsible for the accuracy of any statement or representation made or the consequences arising from the use of or alteration to any information contained in it. These procedures are intended solely as a general resource for practising professionals in the field, operating in the UK, and specialist advice should be obtained where necessary. If you make any changes to this publication, it must be made clear where changes have been made to the original document. The Health Protection Agency (HPA) should at all times be acknowledged.

The HPA is an independent organisation dedicated to protecting people's health. It brings together the expertise formerly in a number of official organisations. More information about the HPA can be found at www.hpa.org.uk.

The HPA aims to be a fully Caldicott compliant organisation. It seeks to take every possible precaution to prevent unauthorised disclosure of patient details and to ensure that patient-related records are kept under secure conditions¹.

More details can be found on the website at www.evaluations-standards.org.uk. Contributions to the development of the documents can be made by contacting standards@hpa.org.uk.

Please note the references are now formatted using Reference Manager software. If you alter or delete text without Reference Manager installed on your computer, the references will not be updated automatically.

Suggested citation for this document:

Health Protection Agency (2004). *Investigation of gastric aspirates and infection screen swabs from neonates*. National Standard Method BSOP 23 Issue 4. http://www.hpa-standardmethods.org.uk/pdf_bacteriology.asp.

INVESTIGATION OF GASTRIC ASPIRATES AND INFECTION SCREEN SWABS FROM NEONATES

Issue no: 4.1 Issue date: 03.05.05 Issued by: Standards Unit, Evaluations and Standards Laboratory Page 2 of 19

Reference no: BSOP 23i4.1

This SOP should be used in conjunction with the series of other SOPs from the Health Protection Agency

www.evaluations-standards.org.uk

Email: standards@hpa.org.uk

INDEX

STATUS OF NATIONAL STANDARD METHODS.....	2
INDEX.....	3
AMENDMENT PROCEDURE.....	4
SCOPE OF DOCUMENT.....	5
INTRODUCTION.....	5
1.0 SAFETY CONSIDERATIONS.....	10
1.1 SPECIMEN COLLECTION.....	10
1.2 SPECIMEN TRANSPORT AND STORAGE.....	10
1.3 SPECIMEN PROCESSING.....	10
2.0 SPECIMEN COLLECTION.....	10
2.1 OPTIMAL TIME OF SPECIMEN COLLECTION.....	10
2.2 CORRECT SPECIMEN TYPE AND METHOD OF COLLECTION.....	10
2.3 ADEQUATE QUANTITY AND APPROPRIATE NUMBER OF SPECIMENS.....	10
3.0 SPECIMEN TRANSPORT AND STORAGE.....	10
3.1 TIME BETWEEN SPECIMEN COLLECTION AND PROCESSING.....	10
3.2 SPECIAL CONSIDERATIONS TO MINIMISE DETERIORATION.....	11
4.0 SPECIMEN PROCESSING.....	11
4.1 TEST SELECTION.....	11
4.2 APPEARANCE.....	11
4.3 MICROSCOPY.....	11
4.4 CULTURE AND INVESTIGATION.....	11
4.5 IDENTIFICATION.....	12
4.6 ANTIBIOTIC SUSCEPTIBILITY TESTING.....	12
REPORTING PROCEDURE.....	13
4.7 MICROSCOPY.....	13
4.8 CULTURE.....	13
4.9 ANTIBIOTIC SUSCEPTIBILITY TESTING.....	13
5.0 REPORTING TO THE HPA (LOCAL AND REGIONAL SERVICES AND CDSC CENTRE FOR INFECTIONS).....	13
REFERENCES.....	15

AMENDMENT PROCEDURE

Controlled document reference	BSOP 23
Controlled document title	Standard Operating Procedure for the investigation of gastric aspirates and infection screen swabs from neonates

Each National Standard Method has an individual record of amendments. The current amendments are listed on this page. The amendment history is available from standards@hpa.org.uk.

On issue of revised or new pages each controlled document should be updated by the copyholder in the laboratory.

Amendment Number/ Date	Issue no. Discarded	Insert Issue no.	Page	Section(s) involved	Amendment
5/ 03.05.05	4	4.1	1	Front page	Redesigned
			2	Status of document	Reworded
			4	Amendment page	Redesigned

INVESTIGATION OF GASTRIC ASPIRATES AND INFECTION SCREEN SWABS FROM NEONATES

Issue no: 4.1 Issue date: 03.05.05 Issued by: Standards Unit, Evaluations and Standards Laboratory Page 4 of 19
Reference no: BSOP 23i4.1

This SOP should be used in conjunction with the series of other SOPs from the Health Protection Agency

www.evaluations-standards.org.uk

Email: standards@hpa.org.uk

STANDARD OPERATING PROCEDURE FOR THE INVESTIGATION OF GASTRIC ASPIRATES AND INFECTION SCREEN SWABS FROM NEONATES

Types of specimens: Gastric aspirates and swabs from ear canal and superficial sites

SCOPE OF DOCUMENT

This SOP describes the processing and bacteriological investigation of gastric aspirates and infection screen swabs from neonates.

INTRODUCTION

Neonatal infection refers to infection occurring during the first four weeks of life.

Infection may be superficial and localised (eg conjunctivitis, superficial skin infections) or deep. Deep infections such as pneumonia, meningitis, septic arthritis or osteomyelitis are frequently systemic, with associated bacteraemia.

The incidence of infection increases with low birth weight or prematurity and may be divided²:

- **early onset** - this occurs in the first 48 hours of life and is usually caused by infection ascending from the maternal genital tract or, less commonly, via the placenta³
- **late-onset** - this occurs after the first 48 hours of life and the organisms may be acquired from the external (eg hospital) environment. Organisms initially colonise superficial sites and the upper respiratory tract and progress to cause widespread sepsis

Infants normally become colonised within the first day of life with flora acquired from the mother and the immediate environment. Coagulase-negative staphylococci predominate on the skin and α -haemolytic streptococci in the upper respiratory tract. In the gut, *Bifidobacterium* species predominate in breast-fed infants, whereas other anaerobes and Enterobacteriaceae are found in greater numbers in bottle-fed babies.

A different pattern of colonisation occurs in infants admitted to a neonatal intensive care unit (NICU) shortly after birth. Colonisation of gut, respiratory tract and skin is delayed, often beyond the third day⁴. The use of antimicrobials and the NICU environment affects the nature of the colonising flora. There is a predominance of organisms such as Gram-negative bacilli and *Staphylococcus aureus*.

Risk factors contributing to the development of neonatal sepsis include⁵:

- congenital abnormalities
- low birth weight
- premature birth
- prolonged rupture of membranes
- maternal fever
- protracted, difficult births
- respiratory distress syndrome
- toxæmia of pregnancy
- invasive procedures and devices (eg intravascular cannulae, shunts, prolonged artificial ventilation)
- infants of mothers who have already had one baby affected by Lancefield group B streptococcal (GBS) sepsis are at increased risk of GBS infection, as is the twin of an affected infant.

Some workers have recommended that infants at special risk of infection and colonised with pathogenic organisms should be treated with antibiotics routinely⁶.

INVESTIGATION OF GASTRIC ASPIRATES AND INFECTION SCREEN SWABS FROM NEONATES

Issue no: 4.1 Issue date: 03.05.05 Issued by: Standards Unit, Evaluations and Standards Laboratory Page 5 of 19

Reference no: BSOP 23i4.1

This SOP should be used in conjunction with the series of other SOPs from the Health Protection Agency

www.evaluations-standards.org.uk

Email: standards@hpa.org.uk

Prolonged rupture of membranes (PROM) is a predisposing factor to both fetal and neonatal infection. Organisms ascending the maternal genital tract may cause intrauterine infection (**amnionitis**). The mother may appear well despite infection of the amniotic fluid and membranes, placenta or fetus. There is increasing evidence that ascending infection is an important cause of preterm labour⁷. Attempts to prevent or control the onset of preterm labour by the increased use of antibiotics^{8,9} may necessitate significant alterations to the methods used for the detection of neonatal infection, aimed at increasing both the speed and specificity of diagnosis.

A variety of microorganisms have also been isolated from the amniotic fluid, amnion and chorion, even at term and in the presence of intact membranes¹⁰, in particular, those associated with bacterial vaginosis¹¹. However, only about 8% of preterm infants whose mothers have clinical evidence of chorioamnionitis develop septicaemia. Bacteraemia, when it occurs, is often transient¹².

Unfortunately, recent randomised trials examining the use of antibiotics versus placebo in various clinical settings of preterm labour, including women with asymptomatic bacterial vaginosis, have not shown significant advantages for the neonate^{13,14}. Indeed, the use of co-amoxiclav for asymptomatic women with preterm rupture of membranes was associated with a significant increase in suspected or proven necrotising enterocolitis in their infants¹⁵. Erythromycin had some advantages over placebo if rupture of membranes was present, but this was lost in the setting of preterm labour with intact membranes^{16,15}.

The clinical manifestations of neonatal sepsis, whatever the source of infection, are frequently non-specific and include respiratory distress, unstable temperature and cardiovascular depression. A bulging or tense fontanelle may be observed in meningitis.

Neonatal pneumonia of early onset may be acquired transplacentally (rubella, cytomegalovirus, listeriosis), or perinatally (associated with asphyxia or possible aspiration of the amniotic fluid)³ or by aspiration of organisms from the birth canal. Late-onset pneumonia is usually of nosocomial origin (eg resulting from use of prolonged mechanical ventilation), although infections acquired during birth (eg *Chlamydia trachomatis*) may also present as late-onset pneumonia.

Neonatal septicaemia and meningitis are also classified into early- and late-onset and may be difficult to diagnose because of the multiplicity of associated risk factors and clinical manifestations^{17,18}.

Organisms associated with sepsis include^{19-21, 22-26}:

- β -haemolytic streptococci, in particular Lancefield group B streptococci
- Enterobacteriaceae
- *S. aureus*
- coagulase-negative staphylococci
- *Listeria monocytogenes*
- Enterococci
- pseudomonads
- yeasts
- *Ureaplasma urealyticum*
- viruses such as echoviruses, herpes simplex

Anaerobes are an extremely rare cause of neonatal infection. Fulminant anaerobic septicaemia tends to be associated with septic abortion or post operative complications.

Lancefield group B streptococci remain the commonest cause of early-onset neonatal sepsis. There is little evidence to support transplacental infection. Organisms may be derived from the mixture of bacteria in the amniotic fluid, if present, or are acquired from the birth canal. Infection is secondary to the multiplication of organisms in the fetal lung³.

Only a small percentage of infants colonised with these organisms develop early onset disease²⁷. Those which do often have other risk factors such as PROM, low birth weight, prematurity or

INVESTIGATION OF GASTRIC ASPIRATES AND INFECTION SCREEN SWABS FROM NEONATES

Issue no: 4.1 Issue date: 03.05.05 Issued by: Standards Unit, Evaluations and Standards Laboratory Page 6 of 19

Reference no: BSOP 23i4.1

This SOP should be used in conjunction with the series of other SOPs from the Health Protection Agency

www.evaluations-standards.org.uk

Email: standards@hpa.org.uk

congenital abnormalities. Late-onset infections associated with acquisition of the organism from the mother or healthcare personnel may also occur.

Intrapartum (ie during labour and delivery) antibiotics given to high risk mothers with Lancefield group B streptococci may prevent ascending infection and have been shown to be effective in reducing the incidence of subsequent early-onset streptococcal disease in high-prevalence populations²⁸⁻³¹.

Coagulase-negative staphylococci (CNS) are commonly associated with the presence of intravascular devices and are the most frequent cause of late infections in neonatal intensive care units (NICUs). They are the commonest organisms isolated from blood cultures in infants in NICUs. Although the technical difficulties inherent in collecting blood cultures aseptically from small infants means that contamination is always possible, the presence of CNS must always be interpreted in the light of the clinical status of the infant at the time blood cultures were collected and its subsequent progress. Infections with CNS may have a more insidious presentation than those caused by Gram-negative bacilli.

It is not possible to interpret the significance of CNS found in superficial swabs and gastric aspirates. The bacterial factors associated with invasion are not well understood. Genotyping studies of CNS colonising and causing infection in NICU infants, while clarifying the molecular epidemiology of the NICU CNS population, cannot predict whether colonised infants will develop infection³².

Gram-negative bacilli are an important cause of early and late-onset neonatal sepsis.

Escherichia coli capsular type K1 is the strain most commonly associated with early onset sepsis³³⁻³⁵. In the NICU setting, it is likely that nosocomial acquisition of hospital strains of *E. coli* occurs, rather than the natural acquisition by neonates of maternal strains of *E. coli*³⁶.

Several outbreaks caused by multi-resistant strains of Enterobacteriaceae and pseudomonads have been reported³⁷⁻⁴⁰. Nosocomial outbreaks of meningitis⁴¹ and enteritis⁴² caused by *Campylobacter jejuni* have also been described.

L. monocytogenes is acquired through transplacental haematogenous spread from bacteraemic mothers or during passage through an infected birth canal. Neonatal listeriosis can present as early-onset or late-onset disease⁴³.

Listeriosis should be considered if there has been maternal febrile illness, or another clinical pointer. Amniotic fluid, placental tissues, maternal blood and vaginal secretions, neonatal blood and CSF should be cultured in addition to the routine infection screening swabs. *L. monocytogenes* DNA can be specifically detected in CSF and tissue by the polymerase chain reaction (PCR)⁴³. Serology is of little use because of cross-reactivity with other Gram-positive bacteria, and culture-positive patients often have no detectable antibodies⁴⁴.

The incidence of neonatal sepsis caused by *L. monocytogenes* has fallen considerably in the UK since a peak in the late 1980s due to public health education of food producers and dietary advice issued to pregnant women⁴⁵.

Other causes of late-onset sepsis include *S. aureus* and enterococci. Systemic fungal infection, particularly due to *Malassezia furfur* as a result of the use of lipid-rich total parenteral nutrition, is also increasing⁴⁶⁻⁴⁸. Infections with *M. furfur* associated with intravascular catheters have also been described⁴⁹. Candida infections are also occurring more frequently⁵⁰.

INVESTIGATION OF NEONATAL SEPSIS. This is a controversial subject.

Isolation of organisms from a significant focus such as blood, CSF, skin vesicle fluid or lesion, or a properly-obtained urine (suprapubic aspirate) remains the most valid method of diagnosing systemic bacterial infection⁵¹.

INVESTIGATION OF GASTRIC ASPIRATES AND INFECTION SCREEN SWABS FROM NEONATES

Issue no: 4.1 Issue date: 03.05.05 Issued by: Standards Unit, Evaluations and Standards Laboratory Page 7 of 19

Reference no: BSOP 23i4.1

This SOP should be used in conjunction with the series of other SOPs from the Health Protection Agency

www.evaluations-standards.org.uk

Email: standards@hpa.org.uk

Although organisms isolated from superficial sites, gastric aspirate and amniotic fluid indicate colonization and may include pathogens responsible for disease, they do not establish the presence of active systemic infection.

Surveillance screening is performed routinely in many neonatal units and may be used to monitor trends in resistant flora and define antibiotic policies. It is seen by some to be useful for predicting the occurrence and defining the aetiology of early- and late-onset neonatal sepsis^{52,53}, but by others as non-contributory or misleading^{54,55}.

An extensive study⁵⁶ of material from the ear canal, nasopharynx, axilla, umbilicus, groin, rectum, stomach and endotracheal tube showed that isolates were rarely the same as those recovered from the blood or fluids of the cerebrospinal, joint, pleural, pericardial and peritoneal spaces. This and other studies have concluded that surface cultures are of limited value in predicting the aetiology of sepsis in the newborn^{57,58}. Thus, in many units this has now been discontinued and restricted to babies who are clinically unwell.

The external ear canal is regarded as the best site to sample in the newborn with known risk factors because it may contain residual amniotic fluid. It has been shown that the ear canal is consistently culture positive in cases of early-onset GBS sepsis whereas other sites are not⁵⁹.

Gastric aspirate flora in the newborn reflects those of the amniotic fluid or birth canal⁶⁰⁻⁶². Gram-stained smears of gastric aspirates may show the presence of polymorphonuclear leucocytes (PMNs), but these may be maternal in origin and may not necessarily represent a fetal inflammatory response⁶³. Their presence is often wrongly interpreted³. However, there are contradictory assertions found in the references and many authorities continue to find their presence significant^{64,65}.

Nasopharyngeal and endotracheal aspirates, together with an external ear canal swab in the newborn have been reported as useful indicators of early sepsis⁶⁶. They are often sampled routinely (as described in BSOP 8) in ventilated babies on NICUs as indicators of the likelihood or actuality of late-onset infection, although the value of this procedure has been widely disputed^{67,68,69}. After the first few hours of life, such specimens do not help to predict the cause of systemic sepsis⁷⁰. However, they may be justified for the purposes of infection control surveillance.

Other diagnostic tests, not described in this SOP, which have been used to try to predict which patients will develop sepsis include:

- An immature to total neutrophil (I:T) ratio⁷¹ (ie a left shift of the PMNs)
- C-reactive protein (CRP)¹⁷. The rise in CRP does not occur early enough in the process of sepsis to be a useful predictor, but persistently normal serial CRP measurements indicate sepsis is an unlikely cause of deterioration⁷²
- Latex agglutination has been used for the detection of Lancefield group B streptococcal antigen in gastric aspirates^{17,73} and concentrated urine¹⁷. Doubts have been cast on the utility of antigen detection as a diagnostic test because of both false-positive and false-negative results⁷⁴

Localised sepsis is investigated by examination of swabs or aspirates of the particular site involved (see relevant SOP).

Eye swabs - neonatal conjunctivitis (ophthalmia neonatorum) is acquired during passage through a birth canal which is infected with *Neisseria gonorrhoeae* or *Chlamydia trachomatis* and is characterised by a sticky discharge. Infection usually becomes apparent between 2-5 days (*N. gonorrhoeae*) or 5-14 days (*C. trachomatis*). Diagnosis is made by microscopy and culture of the discharge or by detection of chlamydial antigen (see BSOP 2).

About one half of all cases of neonatal *C. trachomatis* eye infection subsequently develop pneumonia³³. This usually presents at between 4 and 11 weeks³³.

INVESTIGATION OF GASTRIC ASPIRATES AND INFECTION SCREEN SWABS FROM NEONATES

Issue no: 4.1 Issue date: 03.05.05 Issued by: Standards Unit, Evaluations and Standards Laboratory Page 8 of 19

Reference no: BSOP 23i4.1

This SOP should be used in conjunction with the series of other SOPs from the Health Protection Agency

www.evaluations-standards.org.uk

Email: standards@hpa.org.uk

S. aureus is also a common cause of sticky eye, and in NICUs Enterobacteriaceae are not uncommon causes. *Pseudomonas aeruginosa* conjunctivitis requires aggressive treatment because devastating infection with panophthalmitis, meningitis and bacteraemia may ensue^{75,76}.

UNDER REVIEW

INVESTIGATION OF GASTRIC ASPIRATES AND INFECTION SCREEN SWABS FROM NEONATES

Issue no: 4.1 Issue date: 03.05.05 Issued by: Standards Unit, Evaluations and Standards Laboratory Page 9 of 19

Reference no: BSOP 23i4.1

This SOP should be used in conjunction with the series of other SOPs from the Health Protection Agency

www.evaluations-standards.org.uk

Email: standards@hpa.org.uk

1.0 SAFETY CONSIDERATIONS⁷⁷⁻⁸²

1.1 SPECIMEN COLLECTION

N/A

1.2 SPECIMEN TRANSPORT AND STORAGE

1.2.1. Gastric aspirates

Sterile leakproof container in a sealed plastic bag

1.2.2. Swabs

Sealed plastic bag

1.3 SPECIMEN PROCESSING

Containment Level 2

Laboratory procedures that give rise to infectious aerosols must be conducted in a microbiological safety cabinet

Refer to current guidance on the safe handling of all organisms documented in this SOP

The above guidance should be supplemented with local COSHH and risk assessments

Compliance with postal and transport regulations is essential

2.0 SPECIMEN COLLECTION

2.1 OPTIMAL TIME OF SPECIMEN COLLECTION

Before antimicrobial therapy where possible

Gastric aspirate specimens from newborns should be collected ≤ 4 h post delivery and before feeding

2.2 CORRECT SPECIMEN TYPE AND METHOD OF COLLECTION

Gastric aspirates should be collected according to local protocols

2.3 ADEQUATE QUANTITY AND APPROPRIATE NUMBER OF SPECIMENS

N/A

3.0 SPECIMEN TRANSPORT AND STORAGE

3.1 TIME BETWEEN SPECIMEN COLLECTION AND PROCESSING

Specimens should be transported and processed as soon as possible

3.2 SPECIAL CONSIDERATIONS TO MINIMISE DETERIORATION

Swabs should be transported in Amies transport medium with charcoal⁸³

If processing is delayed, refrigeration is preferable to storage at ambient temperature. Delays of over 48h are undesirable

4.0 SPECIMEN PROCESSING

4.1 TEST SELECTION

N/A

4.2 APPEARANCE

N/A

4.3 MICROSCOPY

4.3.1. Standard

N/A

4.3.2. Supplementary

Gastric aspirates

Select a representative portion of specimen with a sterile loop and make a thin smear on a clean microscope slide for Gram staining

Swabs

Prepare a thin smear on a clean microscope slide for Gram staining

4.4 CULTURE AND INVESTIGATION

4.4.1. Pre-treatment

N/A

4.4.2. Specimen processing

Gastric aspirates

Select a representative portion of specimen with a sterile loop and inoculate a loopful on each agar plate

Swabs

Inoculate each agar plate with swab (see BSOP 54)

For the isolation of individual colonies, spread inoculum with a sterile loop

4.4.3 Culture media, conditions and organisms

For all specimens:

Clinical details/ conditions	Standard media	Incubation			Cultures read	Target organism(s)
		Temp °C	Atmos	Time		
Prolonged rupture of membranes Possible sepsis	Blood agar	35-37	5-10% CO ₂	40-48h	daily	Any of the following: β-haemolytic streptococci Coagulase-negative staphylococci Enterococci Enterobacteriaceae <i>L. monocytogenes</i> Pseudomonads <i>S. aureus</i> Yeasts Pure growth of any other organism
	CLED agar	35-37	air	16-24h	≥16h	

Other organisms for consideration - *N. gonorrhoeae*, *C. trachomatis*, *Mycoplasma* species and *Ureaplasma urealyticum*

4.5 IDENTIFICATION

4.5.1. Minimum level in the laboratory

β-haemolytic streptococci	Lancefield group level
Coagulase-negative staphylococci	"coagulase-negative" level
Enterobacteriaceae	"coliforms" level
Enterococci	genus level
<i>Listeria</i> spp.	species level
<i>Neisseria</i> spp.	species level
Pseudomonads	"pseudomonads" level
<i>S. aureus</i>	species level
Yeasts	"yeasts" level

Organisms may be further identified if clinically or epidemiologically indicated

4.5.2. Referral to Reference Laboratories

Listeria species for typing

Fungi requiring identification and/or susceptibility testing

Organisms with unusual or unexpected resistance, and whenever there is a laboratory or clinical problem or anomaly that requires elucidation

4.6 ANTIBIOTIC SUSCEPTIBILITY TESTING

Refer to SOP on Susceptibility testing (BSOP 45)

INVESTIGATION OF GASTRIC ASPIRATES AND INFECTION SCREEN SWABS FROM NEONATES

Issue no: 4.1 Issue date: 03.05.05 Issued by: Standards Unit, Evaluations and Standards Laboratory Page 12 of 19
Reference no: BSOP 23i4.1

This SOP should be used in conjunction with the series of other SOPs from the Health Protection Agency

www.evaluations-standards.org.uk

Email: standards@hpa.org.uk

REPORTING PROCEDURE

4.7 MICROSCOPY

Gram stain (if performed)

Gastric aspirates and/or ear canal swab - report on WBCs and organisms seen

4.7.1. Microscopy reporting time

Urgent microscopy results to be telephoned or sent electronically

Written report, 16 - 72h

Cases of known or suspected sepsis or amnionitis should be brought to the attention of the medical microbiologist as soon as possible

4.8 CULTURE

Report clinically significant organisms isolated **or**

Report all other growth and include an appropriate comment **or**

Report absence of growth

Also, report results of supplementary investigations

4.8.1. Culture reporting time

Clinically urgent culture results to be telephoned or sent electronically

Written report, 16 - 72h stating, if appropriate, that a further report will be issued

4.9 ANTIBIOTIC SUSCEPTIBILITY TESTING

Report susceptibilities as clinically indicated

5.0 REPORTING TO THE HPA⁸⁴ (LOCAL AND REGIONAL SERVICES AND CDSC CENTRE FOR INFECTIONS)

Refer to the following:

Individual SOPs on organism identification

Health Protection Agency publications

"Reporting to the CDR : A guide for laboratories"

"Hospital infection control : Guidance on the control of infection in hospitals"

Refer to current guidelines on CDSC and COSURV reporting

Local guidelines

Report all the following isolates and clinical conditions:

All cases of ophthalmia neonatorum, *Listeria* species and *N. gonorrhoeae*

UNDER REVIEW

INVESTIGATION OF GASTRIC ASPIRATES AND INFECTION SCREEN SWABS FROM NEONATES

Issue no: 4.1 Issue date: 03.05.05 Issued by: Standards Unit, Evaluations and Standards Laboratory Page 14 of 19

Reference no: BSOP 23i4.1

This SOP should be used in conjunction with the series of other SOPs from the Health Protection Agency

www.evaluations-standards.org.uk

Email: standards@hpa.org.uk

REFERENCES

1. Department of Health NHS Executive: The Caldicott Committee. Report on the review of patient-identifiable information. London. December 1997.
2. Isaacs D, Wilkinson AR, Moxon ER. Surveillance of colonization and late-onset septicaemia in neonates. *J Hosp Infect* 1987;10:114-9.
3. Isaacs D, Moxon ER, editors. Neonatal Infections. Oxford: Butterworth-Heinmann; 1991. p. 1-12
4. Goldmann DA, Leclair J, Macone A. Bacterial colonization of neonates admitted to an intensive care environment. *J Pediatr* 1978;93:288-93.
5. Mims LC, Medawar MS, Perkins JR, Grubb WR. Predicting neonatal infections by evaluation of the gastric aspirate: a study in two hundred and seven patients. *Am J Obstet Gynecol* 1972;114:232-8.
6. MacGregor RR, III, Tunnessen WW, Jr. The incidence of pathogenic organisms in the normal flora of the neonate's external ear and nasopharynx. *Clinical Pediatrics* 1973;12:697-700.
7. Gibbs RS, Romero R, Hillier SL, Eschenbach DA, Sweet RL. A review of premature birth and subclinical infection. *Am J Obstet Gynecol* 1992;166:1515-28.
8. Hauth JC, Goldenberg RL, Andrews WW, DuBard MB, Copper RL. Reduced incidence of preterm delivery with metronidazole and erythromycin in women with bacterial vaginosis. *N Engl J Med* 1995;333:1732-6.
9. Mercer BM, Arheart KL. Antimicrobial therapy in expectant management of preterm premature rupture of the membranes. *Lancet* 1995;346:1271-9.
10. Cassell G, Hauth J, Andrews W, Cutter G, Goldenberg R. Chorioamnion colonization: correlation with gestational age in women delivered following spontaneous labor versus indicated delivery. *Am J Obstet Gynecol* 1993;168:425.
11. Hay PE, Lamont RF, Taylor-Robinson D, Morgan DJ, Ison C, Pearson J. Abnormal bacterial colonisation of the genital tract and subsequent preterm delivery and late miscarriage. *BMJ* 1994;308:295-8.
12. Gilbert GL, editor. Infectious Disease in Pregnancy and the Newborn Infant. Monographs in Clinical Pediatrics ed. Vol 2. Chur: Harwood Academic Publishers; 1991. p. 490-1
13. Kenyon SL, Taylor DJ, Tarnow-Mordi W. Broad-spectrum antibiotics for spontaneous preterm labour: the ORACLE II randomised trial. ORACLE Collaborative Group. *Lancet* 2001;357:989-94.
14. Carey JC, Klebanoff MA, Hauth JC, Hillier SL, Thom EA, Ernest JM, et al. Metronidazole to prevent preterm delivery in pregnant women with asymptomatic bacterial vaginosis. National Institute of Child Health and Human Development Network of Maternal-Fetal Medicine Units. *N Engl J Med* 2000;342:534-40.
15. Kenyon SL, Taylor DJ, Tarnow-Mordi W, for the ORACLE Collaborative Group. Broad-spectrum antibiotics for preterm, prelabour rupture of fetal membranes: the ORACLE I randomised trial. *Lancet* 2001;357:979-88.
16. Kenyon SL, Taylor DJ, Tarnow-Mordi W. Broad-spectrum antibiotics for spontaneous preterm labour: the ORACLE II randomised trial. ORACLE Collaborative Group. *Lancet* 2001;357:989-94.

INVESTIGATION OF GASTRIC ASPIRATES AND INFECTION SCREEN SWABS FROM NEONATES

Issue no: 4.1 Issue date: 03.05.05 Issued by: Standards Unit, Evaluations and Standards Laboratory Page 15 of 19

Reference no: BSOP 23i4.1

This SOP should be used in conjunction with the series of other SOPs from the Health Protection Agency

www.evaluations-standards.org.uk

Email: standards@hpa.org.uk

17. Philip AGS. Diagnostic tests for bacterial infection in the newborn. In: de Louvois J, Harvey D, editors. *Infection in the Newborn*. Vol 6. Chichester: John Wiley & Sons; 1990. p. 49-59.
18. Philip AGS, editor. *Neonatal sepsis and meningitis*. Boston: GK Hall & Co; 1985. p. 1-5
19. Goldmann DA, Leclair J, Macone A. Bacterial colonization of neonates admitted to an intensive care environment. *J Pediatr* 1978;93:288-93.
20. Mims LC, Medawar MS, Perkins JR, Grubb WR. Predicting neonatal infections by evaluation of the gastric aspirate: a study in two hundred and seven patients. *Am J Obstet Gynecol* 1972;114:232-8.
21. MacGregor RR, III, Tunnessen WW, Jr. The incidence of pathogenic organisms in the normal flora of the neonate's external ear and nasopharynx. *Clinical Pediatrics* 1973;12:697-700.
22. Sprunt K, Leidy G, Redman W. Abnormal colonization of neonates in an intensive care unit: means of identifying neonates at risk of infection. *Pediatr Res* 1978;12:998-1002.
23. Wilson MG, Armstrong DH, Nelson RC, Boak RA. Prolonged rupture of fetal membranes. *Am J Dis Child* 1964;107:138-46.
24. Evans ME, Schaffner W, Federspiel CF, Cotton RB, McKee KT, Jr., Stratton CW. Sensitivity, specificity, and predictive value of body surface cultures in a neonatal intensive care unit. *JAMA* 1988;259:248-52.
25. Hemming VG, Overall JC, Jr., Britt MR. Nosocomial infections in a newborn intensive-care unit. Results of forty-one months of surveillance. *N Engl J Med* 1976;294:1310-6.
26. Davies PA. Bacterial infection in the fetus and newborn. *Arch Dis Child* 1971;46:1-27.
27. Berner R. Group B streptococci during pregnancy and infancy. [Review] [57 refs]. *Current Opinion in Infectious Diseases* 2002;15:307-13.
28. Boyer KM, Gadzala CA, Kelly PD, Gotoff SP. Selective intrapartum chemoprophylaxis of neonatal group B streptococcal early-onset disease. III. Interruption of mother-to- infant transmission. *J Infect Dis* 1983;148:810-6.
29. Boyer KM, Gotoff SP. Prevention of early-onset neonatal group B streptococcal disease with selective intrapartum chemoprophylaxis. *N Engl J Med* 1986;314:1665-9.
30. Schrag SJ, Zywicki S, Farley MM, Reingold AL, Harrison LH, Lefkowitz LB, et al. Group B streptococcal disease in the era of intrapartum antibiotic prophylaxis. *N Engl J Med* 2000;342:15-20.
31. Interim "good practice" recommendations for the prevention of early onset neonatal group B streptococcal (GBS) infection in UK. http://www.hpa.org.uk/infections/topics_az/strepto/guidelines.htm.
32. de Silva GD, Justice A, Wilkinson AR, Buttery J, Herbert M, Day NP, et al. Genetic population structure of coagulase-negative staphylococci associated with carriage and disease in preterm infants. *Clin Infect Dis* 2001;33:1520-8.
33. Saez-Llorens X, McCracken GH J. Perinatal bacterial diseases. In: Feigin RD, Cherry JD, editors. *Textbook of Pediatric Infectious Diseases*. 4th ed. Vol 1. Philadelphia: WB Saunders Company; 1998. p. 892-926.

INVESTIGATION OF GASTRIC ASPIRATES AND INFECTION SCREEN SWABS FROM NEONATES

Issue no: 4.1 Issue date: 03.05.05 Issued by: Standards Unit, Evaluations and Standards Laboratory Page 16 of 19

Reference no: BSOP 23i4.1

This SOP should be used in conjunction with the series of other SOPs from the Health Protection Agency

www.evaluations-standards.org.uk

Email: standards@hpa.org.uk

34. Murono K, Fujita K, Yoshikawa M, Saijo M, Inyaku F, Kakehashi H, et al. Acquisition of non-maternal Enterobacteriaceae by infants delivered in hospitals. *J Pediatr* 1993;122:120-5.
35. Sarff LD, McCracken GH, Schiffer MS, Glode MP, Robbins JB, Orskov I, et al. Epidemiology of *Escherichia coli* K1 in healthy and diseased newborns. *Lancet* 1975;1:1099-104.
36. Fujita K, Murono K. Nosocomial acquisition of *Escherichia coli* by infants delivered in hospitals. [Review] [23 refs]. *Journal of Hospital Infection* 1996;32:277-81.
37. Becks VE, Lorenzoni NM. *Pseudomonas aeruginosa* outbreak in a neonatal intensive care unit: a possible link to contaminated hand lotion. *Am J Infect Control* 1995;23:396-8.
38. Garland SM, Mackay S, Tabrizi S, Jacobs S. *Pseudomonas aeruginosa* outbreak associated with a contaminated blood- gas analyser in a neonatal intensive care unit. *J Hosp Infect* 1996;33:145-51.
39. Acolet D, Ahmet Z, Houang E, Hurley R, Kaufmann ME. *Enterobacter cloacae* in a neonatal intensive care unit: account of an outbreak and its relationship to use of third generation cephalosporins. *J Hosp Infect* 1994;28:273-86.
40. Umasankar S, Mridha EU, Hannan MM, Fry CM, Azadian BS. An outbreak of *Salmonella enteritidis* in a maternity and neonatal intensive care unit. *J Hosp Infect* 1996;34:117-22.
41. Goossens H, Henocque G, Kremp L, Rocque J, Boury R, Alanio G, et al. Nosocomial outbreak of *Campylobacter jejuni* meningitis in newborn infants. *Lancet* 1986;2:146-9.
42. Terrier A, Altwegg M, Bader P, von Graevenitz A. Hospital epidemic of neonatal *Campylobacter jejuni* infection. *Lancet* 1985;2:1182.
43. Broome C, Pinner R, Schuchat A. *Listeria monocytogenes*. In: Gorbach SL, Bartlett JG, Blacklow NR, editors. *Infectious Diseases*. 2nd ed. Philadelphia: WB Saunders Company; 1998. p. 1750-5.
44. Schuchat A, Swaminathan B, Broome CV. Epidemiology of human listeriosis. *Clin Microbiol Rev* 1991;4:169-83.
45. Smerdon WJ, Jones R, McLauchlin J, Reacher M. Surveillance of listeriosis in England and Wales, 1995-1999. *Commun Dis Public Health* 2001;4:188-93.
46. Papavassilis C, Mach KK, Mayser PA. Medium-chain triglycerides inhibit growth of *Malassezia*: implications for prevention of systemic infection. *Crit Care Med* 1999;27:1781-6.
47. Kim EH, Cohen RS, Ramachandran P, Glasscock GF. Adhesion of percutaneously inserted Silastic central venous lines to the vein wall associated with *Malassezia furfur* infection. *JPEN J Parenter Enteral Nutr* 1993;17:458-60.
48. Weiss SJ, Schoch PE, Cunha BA. *Malassezia furfur* fungemia associated with central venous catheter lipid emulsion infusion. *Heart Lung* 1991;20:87-90.
49. Isaacs D, Moxon ER, editors. *Neonatal Infections*. Oxford: Butterworth-Heinemann; 1991. p. 140-8
50. Dankner WM, Spector SA, Fierer J, Davis CE. *Malassezia* fungemia in neonates and adults: complication of hyperalimentation. *Rev Infect Dis* 1987;9:743-53.
51. Powell KR, Marcy SM. Laboratory aids for diagnosis of neonatal sepsis. In: Remington JS, Klein JO, editors. *Infectious Diseases of the Fetus and Newborn Infant*. 4th ed. Philadelphia: WB Saunders Company; 1995. p. 1223-40.

INVESTIGATION OF GASTRIC ASPIRATES AND INFECTION SCREEN SWABS FROM NEONATES

Issue no: 4.1 Issue date: 03.05.05 Issued by: Standards Unit, Evaluations and Standards Laboratory Page 17 of 19

Reference no: BSOP 23i4.1

This SOP should be used in conjunction with the series of other SOPs from the Health Protection Agency

www.evaluations-standards.org.uk

Email: standards@hpa.org.uk

52. Finelli L, Livengood JR, Saiman L. Surveillance of pharyngeal colonization: detection and control of serious bacterial illness in low birth weight infants. *Pediatr Infect Dis J* 1994;13:854-9.
53. Lee PY, Holliman RE, Davies EG. Surveillance cultures on neonatal intensive care units. *J Hosp Infect* 1995;29:233-6.
54. Goossens H, Henocque G, Kremp L, Rocque J, Boury R, Alanio G, et al. Nosocomial outbreak of *Campylobacter jejuni* meningitis in newborn infants. *Lancet* 1986;2:146-9.
55. Webber S, Wilkinson AR, Lindsell D, Hope PL, Dobson SR, Isaacs D. Neonatal pneumonia. *Arch Dis Child* 1990;65:207-11.
56. Evans ME, Schaffner W, Federspiel CF, Cotton RB, McKee KT, Jr., Stratton CW. Sensitivity, specificity, and predictive value of body surface cultures in a neonatal intensive care unit. *JAMA* 1988;259:248-52.
57. Evans ME, Schaffner W, Federspiel CF, Cotton RB, McKee KT, Jr., Stratton CW. Sensitivity, specificity, and predictive value of body surface cultures in a neonatal intensive care unit. *JAMA* 1988;259:248-52.
58. Fulginiti VA, Ray CG. Body surface cultures in the newborn infant. An exercise in futility, wastefulness, and inappropriate practice. *Am J Dis Child* 1988;142:19-20.
59. Thompson PJ, Greenough A, Gamsu HR, Nicolaidis KH, Philpott-Howard J. Congenital bacterial sepsis in very preterm infants. *J Med Microbiol* 1992;36:117-20.
60. Anderson GS, Green CA, Neligan GA, Newell DJ, Russell JK. Congenital bacterial pneumonia. *Lancet* 1962;2:585-7.
61. Vasan U, Lim DM, Greenstein RM, Raye JR. Origin of gastric aspirate polymorphonuclear leukocytes in infants born after prolonged rupture of membranes. *J Pediatr* 1977;91:69-72.
62. Yeung CY, Tam AS. Gastric aspirate findings in neonatal pneumonia. *Arch Dis Child* 1972;47:735-40.
63. Weiss SJ, Schoch PE, Cunha BA. *Malassezia furfur* fungemia associated with central venous catheter lipid emulsion infusion. *Heart Lung* 1991;20:87-90.
64. Sherman MP, Goetzman BW, Ahlfors CE, Wennberg RP. Tracheal aspiration and its clinical correlates in the diagnosis of congenital pneumonia. *Pediatrics* 1980;65:258-63.
65. Radetsky M. The laboratory evaluation of newborn sepsis. *Current Opinion in Infectious Diseases* 1995;8:191-9.
66. Sherman MP, Goetzman BW, Ahlfors CE, Wennberg RP. Tracheal aspiration and its clinical correlates in the diagnosis of congenital pneumonia. *Pediatrics* 1980;65:258-63.
67. Finelli L, Livengood JR, Saiman L. Surveillance of pharyngeal colonization: detection and control of serious bacterial illness in low birth weight infants. *Pediatr Infect Dis J* 1994;13:854-9.
68. Slagle TA, Bifano EM, Wolf JW, Gross SJ. Routine endotracheal cultures for the prediction of sepsis in ventilated babies. *Arch Dis Child* 1989;64:34-8.
69. Thureen PJ, Moreland S, Rodden DJ, Merenstein GB, Levin M, Rosenberg AA. Failure of tracheal aspirate cultures to define the cause of respiratory deteriorations in neonates. *Pediatr Infect Dis J* 1993;12:560-4.

INVESTIGATION OF GASTRIC ASPIRATES AND INFECTION SCREEN SWABS FROM NEONATES

Issue no: 4.1 Issue date: 03.05.05 Issued by: Standards Unit, Evaluations and Standards Laboratory Page 18 of 19

Reference no: BSOP 23i4.1

This SOP should be used in conjunction with the series of other SOPs from the Health Protection Agency

www.evaluations-standards.org.uk

Email: standards@hpa.org.uk

70. Webber S, Wilkinson AR, Lindsell D, Hope PL, Dobson SR, Isaacs D. Neonatal pneumonia. Arch Dis Child 1990;65:207-11.
71. Manroe BL, Weinberg AG, Rosenfeld CR, Browne R. The neonatal blood count in health and disease. I. Reference values for neutrophilic cells. J Pediatr 1979;95:89-98.
72. Pourcyrous M, Bada HS, Korones SB, Baselski V, Wong SP. Significance of serial C-reactive protein responses in neonatal infection and other disorders. Pediatrics 1993;92:431-5.
73. Poilane I, Adam MN, Torlotin JC, Collignon A. Evaluation of a rapid agglutination test for detection of group B streptococci in the gastric aspirates of neonates. Eur J Clin Microbiol Infect Dis 1995;14:815-7.
74. Williamson M, Fraser SH, Tilse M. Failure of the urinary group B streptococcal antigen test as a screen for neonatal sepsis. Arch Dis Child Fetal Neonatal Ed 1995;73:F109-F111.
75. Lohrer R, Belohradsky BH. Bacterial endophthalmitis in neonates. Eur J Pediatr 1987;146:354-9.
76. Boyle EM, Ainsworth JR, Levin AV, Campbell AN, Watkinson M. Ophthalmic Pseudomonas infection in infancy. Archives of Disease in Childhood: Fetal & Neonatal Edition 2001;85:F139-F140.
77. Advisory Committee on Dangerous Pathogens. Categorisation of biological agents according to hazard and categories of containment. 4th ed. Suffolk: HSE Books; 1995. Supplements 1, 1998 and 2, 2000.
78. Public Health Laboratory Service Standing Advisory Committee on Laboratory Safety. Safety Precautions: Notes for Guidance. 4th ed. London: Public Health Laboratory Service (PHLS); 1993.
79. Control of Substances Hazardous to Health Regulations 2002. General COSHH Approved Code of Practice and Guidance, L5. Suffolk: HSE Books; 2002.
80. Health and Safety Executive. 5 steps to risk assessment: a step by step guide to a safer and healthier workplace, IND (G) 163 (REVL). Suffolk: HSE Books; 2002.
81. Health and Safety Executive. A guide to risk assessment requirements: common provisions in health and safety law, IND (G) 218 (L). Suffolk: HSE Books; 2002.
82. Health Services Advisory Committee. Safety in Health Service laboratories. Safe working and the prevention of infection in clinical laboratories and similar facilities. 2nd ed. Suffolk: HSE Books; 2003.
83. Barber S, Lawson PJ, Grove DI. Evaluation of bacteriological transport swabs. Pathology 1998;30:179-82.
84. PHLS CDSC. Reporting to the PHLS Communicable Disease Surveillance Centre: a reference for laboratories. May. 2001.

INVESTIGATION OF GASTRIC ASPIRATES AND INFECTION SCREEN SWABS FROM NEONATES

Issue no: 4.1 Issue date: 03.05.05 Issued by: Standards Unit, Evaluations and Standards Laboratory Page 19 of 19

Reference no: BSOP 23i4.1

This SOP should be used in conjunction with the series of other SOPs from the Health Protection Agency

www.evaluations-standards.org.uk

Email: standards@hpa.org.uk